

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTALDB1623

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAR 15	WPIDS/WPIX enhanced with new FRAGHITSTR display format
NEWS	3	MAR 16	CASREACT coverage extended
NEWS	4	MAR 20	MARPAT now updated daily
NEWS	5	MAR 22	LWPI reloaded
NEWS	6	MAR 30	RDISCLOSURE reloaded with enhancements
NEWS	7	APR 02	JICST-EPLUS removed from database clusters and STN
NEWS	8	APR 30	GENBANK reloaded and enhanced with Genome Project ID field
NEWS	9	APR 30	CHEMCATS enhanced with 1.2 million new records
NEWS	10	APR 30	CA/CAPplus enhanced with 1870-1889 U.S. patent records
NEWS	11	APR 30	INPADOC replaced by INPADOCDB on STN
NEWS	12	MAY 01	New CAS web site launched
NEWS	13	MAY 08	CA/CAPplus Indian patent publication number format defined
NEWS	14	MAY 14	RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS	15	MAY 21	BIOSIS reloaded and enhanced with archival data
NEWS	16	MAY 21	TOXCENTER enhanced with BIOSIS reload
NEWS	17	MAY 21	CA/CAPplus enhanced with additional kind codes for German patents
NEWS	18	MAY 22	CA/CAPplus enhanced with IPC reclassification in Japanese patents
NEWS	19	JUN 27	CA/CAPplus enhanced with pre-1967 CAS Registry Numbers
NEWS	20	JUN 29	STN Viewer now available
NEWS	21	JUN 29	STN Express, Version 8.2, now available
NEWS	22	JUL 02	LEMBASE coverage updated
NEWS	23	JUL 02	LMEDLINE coverage updated
NEWS	24	JUL 02	SCISEARCH enhanced with complete author names
NEWS	25	JUL 02	CHEMCATS accession numbers revised
NEWS	26	JUL 02	CA/CAPplus enhanced with utility model patents from China
NEWS EXPRESS	29	JUNE 2007:	CURRENT WINDOWS VERSION IS V8.2, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 05 JULY 2007.
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS LOGIN			Welcome Banner and News Items
NEWS IPC8			For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 18:43:07 ON 11 JUL 2007

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CAPLUS' ENTERED AT 18:43:17 ON 11 JUL 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 11 Jul 2007 VOL 147 ISS 3

FILE LAST UPDATED: 10 Jul 2007 (20070710/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> S (POLY (5A) ICLC)

706494 POLY

2 POLIES

706495 POLY

(POLY OR POLIES)

105 ICLC

L1 94 (POLY (5A) ICLC)

=> d scan

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 1-6 (Pharmacology)

TI Cellular regulation by immunomodifiers MVE-2 and poly ICLC and their therapeutic application

ST immunomodifier antitumor cell regulation; MVE 2 antitumor immunomodifier; polymer lysine nucleotide immunomodifier antitumor

IT Lymphokines and Cytokines

RL: BIOL (Biological study)

(colony-stimulating factor secretion by, immunomodifiers effect on, neoplasm inhibition in relation to)

IT Neoplasm inhibitors

(immunomodifiers, cell regulation in relation to)

IT Immune adjuvants

(lysine-nucleotide polymer and MVE-2 as, neoplasm inhibition in relation to)

IT Prostaglandins

RL: FORM (Formation, nonpreparative)

(E, formation of, immunomodifiers effect on, neoplasm inhibition in relation to)

IT Macrophage

(cytotoxic, immunomodifiers, neoplasm inhibition in relation to)

IT Lymphocyte

(natural killer, immunomodifiers, neoplasm inhibition in relation to)

IT 50-18-0

RL: BIOL (Biological study)

(neoplasm inhibition by MVE-2 and, immune modulation in)

IT 27100-68-1 59789-29-6

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(neoplasm inhibition by, immune modulation in, cellular regulation in relation to)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> l1 and prep/rl

4430063 PREP/RL

L2 2 L1 AND PREP/RL

=> d l2 1-2

L2 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2004:18891 CAPLUS

DN 140:71067

TI Method for preparation of large volume batches of poly-ICLC with increased biological potency, and therapeutic, clinical and veterinary uses thereof

IN Salazar, Andres

PA Oncovir, Inc., USA

SO U.S. Pat. Appl. Publ., 14 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004005998	A1	20040108	US 2003-611614	20030701
	WO 2005102278	A1	20051103	WO 2003-US20828	20030701
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW,			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	AU 2003248791	A1	20051109	AU 2003-248791	20030701
	EP 1778186	A1	20070502	EP 2003-819324	20030701
	R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PT, RO, SE, SI, SK, TR			
PRAI	US 2002-393713P	P	20020703		
	WO 2003-US20828	W	20030701		

L2 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1982:135517 CAPLUS

DN 96:135517

TI Polyribonucleosinic-polyribocytidylic acid-poly-L-lysine complex [poly(ICL)] without carboxymethylcellulose (CMC): a new primate-effective interferon inducer

AU Riley, Freddie L.; Morin, Martin L.; Lvovsky, Eduard; Stephens, Edward E.; Levy, Hilton B.

CS Natl. Inst. Allergy Infect. Dis., Bethesda, MD, 21701, USA

SO Proceedings of the Society for Experimental Biology and Medicine (1982), 169(2), 183-8

CODEN: PSEBAA; ISSN: 0037-9727

DT Journal

LA English

=> d his

(FILE 'HOME' ENTERED AT 18:43:07 ON 11 JUL 2007)

FILE 'CAPLUS' ENTERED AT 18:43:17 ON 11 JUL 2007

L1 94 S (POLY (5A) ICLC)  
L2 2 L1 AND PREP/RL

=> d scan l1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 15-5 (Immunochemistry)  
TI Secretion of colony-stimulating factors by human monocytes and bone marrow cells after in vitro treatment with biological response modifiers  
ST interferon monocyte colony stimulating factor  
IT Interferons  
RL: BIOL (Biological study)  
(colony-stimulating factor secretion by monocyte stimulation by, of humans)  
IT Monocyte  
(colony-stimulating factor secretion by, biol. response modifiers stimulation of, of humans)  
IT Hematopoietic precursor cell  
(myeloid, biol. response modifiers effect on, of humans)  
IT 64769-70-6 75985-31-8  
RL: BIOL (Biological study)  
(colony-stimulating factor secretion by bone marrow cells and monocytes stimulation by, of humans)  
IT 62683-29-8  
RL: BIOL (Biological study)  
(secretion of, by bone marrow cells and monocytes, biol. response modifiers stimulation of, of humans)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-5 (Pharmacology)  
TI Use of the antiviral and immune modulator, poly(ICLC), in the treatment of AIDS  
ST polyICLC AIDS antiviral  
IT Lymphocyte  
(function of human, in AIDS, poly(ICLC) effect on)  
IT Immunomodulators  
Virucides and Virustats  
(poly(ICLC) as, AIDS in humans response to)  
IT Acquired immune deficiency syndrome  
(treatment of, by poly(ICLC), in humans)  
IT 30516-87-1, AZT  
RL: BIOL (Biological study)  
(AIDS treatment by poly(ICLC) and, in humans)  
IT 59789-29-6, Poly(ICLC)  
RL: BIOL (Biological study)  
(AIDS treatment by, in humans)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-5 (Pharmacodynamics)  
TI Effect of treatment with exogenous interferon, polyriboinosinic-polyribocytidylic acid, or polyriboinosinic-polyribocytidylic acid-poly-L-lysine complex on Herpesvirus hominis infections in mice  
ST interferon inducer herpesvirus infection; ribonucleotide hypervirus infection; polyriboinosinic acid polyribocytidylic acid herpesvirus;

polyribocytidylate polylysine polyriboninosinate herpesvirus  
 IT Interferons  
 RL: BIOL (Biological study)  
 (herpesvirus infection treatment with)  
 IT Virus, animal  
 (Herpesvirus hominis type 2, infection with, interferon and interferon  
 inducers treatment of)  
 IT 24939-03-5 64769-70-6  
 RL: BIOL (Biological study)  
 (herpesvirus infection treatment with)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 1-7 (Pharmacology)  
 TI Purification of carboxymethylcellulose decreases toxicity of poly  
 ICLC in mice  
 ST polyinosinylcytidylate polylysine carboxymethylcellulose interferon  
 toxicity  
 IT Interferons  
 RL: PRP (Properties)  
 (induction of, with polyinosinic polycytidylic acid-polylysine-  
 carboxymethylcellulose, carboxymethylcellulose purification effect on,  
 toxicity in relation to)  
 IT 59789-29-6  
 RL: PRP (Properties)  
 (toxicity of, carboxymethylcellulose purification decrease of, interferon  
 induction in relation to)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 15-5 (Immunochemistry)  
 TI TNF- $\alpha$  is a principal cytokine involved in the recruitment of NK  
 cells to liver parenchyma  
 ST tumor necrosis factor natural killer lymphocyte; liver NK lymphocyte tumor  
 necrosis factor  
 IT Liver  
 (natural killer lymphocyte adhesion to endothelium of, tumor necrosis  
 factor- $\alpha$  involvement in)  
 IT Adhesion  
 (bio-, of natural killer cells to liver endothelium, tumor necrosis  
 factor- $\alpha$  involvement in)  
 IT Lymphocyte  
 (natural killer cell, adhesion of, to liver endothelium, tumor necrosis  
 factor- $\alpha$  involvement in)  
 IT Lymphokines and Cytokines  
 RL: BIOL (Biological study)  
 (tumor necrosis factor- $\alpha$ , in natural killer cell adhesion to  
 liver endothelium)  
 IT 24939-03-5, Polyinosinic-polycytidylic acid 25104-18-1, Poly-L-lysine  
 RL: BIOL (Biological study)  
 (hepatic natural killer lymphocyte activity enhancement by)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 15-13 (Immunochemistry)  
 Section cross-reference(s): 1  
 TI Adjuvant effects of low doses of a nuclease-resistant derivative of  
 polyinosinic acid-polycytidylic acid on antibody responses of monkeys to  
 inactivated Venezuelan equine encephalomyelitis virus vaccine  
 ST immune adjuvant polyinosinate polycytidylate; polylysine polyIC immune  
 adjuvant; CM cellulose polyIC immune adjuvant  
 IT Immune adjuvants

(polyinosinic acid-polycytidylic acid-polylysine-carboxymethylcellulose complexes as, at low doses)

IT 59789-29-6  
RL: BIOL (Biological study)  
(immune adjuvanticity of, at low doses)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-6 (Pharmacology)  
TI Immunotherapy of metastasis: comparative efficacy of BRMs for the treatment of transplantable and autochthonous tumors  
ST biol response modifier antimetastatic; immunomodulator antimetastatic  
IT Immune adjuvants  
(neoplasm metastasis treatment with)  
IT Neoplasm inhibitors  
(metastasis, immunomodulating agents as)  
IT 27100-68-1 37339-90-5 39325-01-4 59789-29-6 79335-75-4  
RL: BIOL (Biological study)  
(neoplasm metastasis treatment with, immunomodulating activity in)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-5 (Pharmacology)  
TI Killing of Leishmania donovani amastigotes by poly ICLC in hamsters  
ST Leishmania amastigote antileishmanial polyinosinate polycytidylate arginine  
IT Leishmania donovani  
(killing of Leishmania donovani amastigotes by poly ICLC and L-arginine in hamsters)  
IT Microbicidal and microbiostatic action  
(leishmanicidal, killing of Leishmania donovani amastigotes by poly ICLC and L-arginine in hamsters)  
IT 10102-43-9, Nitric oxide, biological studies 125978-95-2, Nitric oxide synthase  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(in killing of Leishmania donovani amastigotes by poly ICLC and L-arginine in hamsters)  
IT 74-79-3, L-Arginine, biological studies 59789-29-6, Poly ICLC  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(killing of Leishmania donovani amastigotes by poly ICLC and L-arginine in hamsters)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 15-13 (Immunochemistry)  
Section cross-reference(s): 1  
TI Immune modulating effects of poly ICLC  
ST immune adjuvant interferon inducer; polynucleotide polylysine complex immune adjuvant  
IT Immune adjuvants  
((poly I-poly C)-polylysine-carboxymethylcellulose complexes as, interferon induction in relation to)  
IT Vaccines  
(immune adjuvant activity of interferon inducers with)  
IT Interferons  
RL: PRP (Properties)  
(induction of, with (poly I-poly C)-polylysine-

carboxymethylcellulose complexes, immune adjuvant activity in relation to)

IT 59789-29-6

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(immune adjuvant activity of, interferon induction in relation to)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 1-6 (Pharmacology)

TI Toxicity of polyinosinic-polycytidylic acid admixed with poly-L-lysine and solubilized with carboxymethylcellulose in mice  
ST poly ICLC toxicity; polyinosinate polycytidylate polylysine toxicity

IT Liver, toxic chemical and physical damage

Lung, toxic chemical and physical damage

(polyinosinate-polycytidylate-polylysine toxicity to)

IT 59789-29-6, Poly (ICLC)

RL: PRP (Properties)

(toxicity of, to liver and lungs)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

TI Immunochemotherapy for Leishmania donovani infection in golden hamsters: combinatorial action of poly ICLC plus L-arginine and sodium stibogluconate (Stibanate)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 1-7 (Pharmacology)

Section cross-reference(s): 15

TI Interferon induction and therapy of brain tumors in rats by poly (ICLC)

ST polyriboinosinate polyribocytidylate analog interferon antitumor; brain tumor polyriboinosinate polyribocytidylate analog; polylysine CM cellulose polynucleotide interferon induction

IT Interferons

RL: PRP (Properties)

(induction of, by poly(I)·poly(C)-polylysine-CM cellulose, in brain tumor immunotherapy)

IT Immune adjuvants

(interferon induction and brain tumor inhibition by poly(I)·poly(C)-polylysine-CM cellulose in relation to)

IT Brain, neoplasm

(interferon induction by poly(I)·poly(C)-polylysine-CM cellulose in immunotherapy of)

IT Neoplasm inhibitors

(poly(I)·poly(C)-polylysine-CM cellulose as, for brain neoplasm, interferon induction in relation to)

IT 59789-29-6

RL: BIOL (Biological study)

(interferon induction and brain tumor inhibition by)

IT 24939-03-5

RL: BIOL (Biological study)

(interferon induction and brain tumor inhibition by analog of)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 15-5 (Immunochemistry)

Section cross-reference(s): 1

TI Activation of natural killer cells in newborn piglets by interferon

induction  
ST natural killer cell interferon activation; piglet immunity interferon inducer  
IT Interferons  
RL: PRP (Properties)  
(natural killer cell activation by exogenous and induction of, in piglets)  
IT Swine  
(natural killer lymphocyte activation in newborn and weanling, interferon induction of)  
IT Lymphocyte  
(natural killer, activation of, by interferon, in piglets)  
IT Virus, animal  
(transmissible gastroenteritis, infection with, in piglets, interferon induction inhibition of)  
IT 59789-29-6, Poly ICLC  
RL: BIOL (Biological study)  
(interferon induction by, natural killer cell response to, in piglets)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
IC ICM A61K038-16  
ICS A61K031-716  
INCL 514002000; 514057000  
CC 1-12 (Pharmacology)  
TI Method for preparation of large volume batches of poly-ICLC with increased biological potency, and therapeutic, clinical and veterinary uses thereof  
ST poly ICLC prodn therapeutic gene regulation; infection  
IT CD antigens  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(CD106; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Enzymes, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(DNA helicase; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Nervous system, disease  
(Guillain-Barre syndrome; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Enzymes, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(RNA helicase; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Infection  
Respiratory system, disease  
(SARS (severe acute respiratory syndrome); preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Cell adhesion molecules  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(VCAM-1 (vascular cell adhesion mol. 1); preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(actin filament-associated; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Neuroglia, neoplasm  
(astrocytoma; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)  
IT Respiratory system, disease  
(bovine respiratory complex; preparation of large volume batches of



poly-ICLC with increased biol. potency, and therapeutic use)

IT Intestine, neoplasm  
(colon; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Enzymes, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (dsRNA-inducible; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Translation initiation factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (eIF-2; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Encephalitis  
(equine viral; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Neuroglia, neoplasm  
(glioblastoma; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Immune disease  
(immune neuropathy; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Interferons  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (induction; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Drug delivery systems  
(injections, i.m.; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Drug delivery systems  
(injections, i.v.; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Ionizing radiation  
(injury; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (interferon regulatory factor; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Infection  
(microbial; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Drug delivery systems  
(nasal; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Astrocyte  
(neoplasm, astrocytoma; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Nerve, disease  
(neuropathy, immune; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Drug delivery systems  
(oral; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (p56; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT AIDS (disease)  
Adenoviridae  
Anti-AIDS agents  
Anti-infective agents  
Antitumor agents  
Antiviral agents

Apoptosis  
 Biological transport  
 Brain, neoplasm  
 Cell cycle  
 Coronavirus  
 Cytoskeleton  
 Dengue virus  
 Drug delivery systems  
 Ebola virus  
 Extracellular matrix  
 Filovirus  
 Flavivirus  
 Foot-and-mouth disease virus  
 Hepatitis virus  
 Herpesviridae  
 Human  
 Human adenovirus  
 Human herpesvirus  
 Human immunodeficiency virus  
 Immunostimulants  
 Influenza virus  
 Japanese encephalitis virus  
 Leukemia  
 Lung, neoplasm  
 Lymphoma  
 Mammary gland, neoplasm  
 Melanoma  
 Metabolism  
 Multiple sclerosis  
 Neuroglia, neoplasm  
 Porcine respiratory and reproductive syndrome virus  
 Poxviridae  
 RNA formation  
 Radioprotectants  
 Sarcoma  
 Signal transduction, biological  
 Translation, genetic  
 Vaccines  
 Vaccinia virus  
 Variola virus  
 West Nile virus  
 Yellow fever virus  
 (preparation of large volume batches of poly-ICLC with  
 increased biol. potency, and therapeutic use)

IT

Cytokines  
 Double stranded RNA  
 Gene, animal  
 Growth factors, animal  
 Tumor necrosis factors  
 p53 (protein)  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (preparation of large volume batches of poly-ICLC with  
 increased biol. potency, and therapeutic use)

IT

Kidney, neoplasm  
 (renal cell carcinoma; preparation of large volume batches of poly-  
 ICLC with increased biol. potency, and therapeutic use)

IT

Carcinoma  
 (renal cell; preparation of large volume batches of poly-  
 ICLC with increased biol. potency, and therapeutic use)

IT

Drug delivery systems  
 (sublingual; preparation of large volume batches of poly-  
 ICLC with increased biol. potency, and therapeutic use)

IT

Drug delivery systems  
 (topical; preparation of large volume batches of poly-ICLC  
 with increased biol. potency, and therapeutic use)

IT Drug delivery systems  
(transdermal; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Drug toxicity  
(vaccine; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Blood vessel, disease  
(vasculitides; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT Infection  
(viral; preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT 62031-54-3, Fibroblast growth factor 69106-44-1, 2',5'-Oligoadenylate synthetase 91608-96-7, p68 Protein kinase 105913-11-9, Plasminogen activator 141907-41-7, Matrix metalloproteinase  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT 59789-29-6P, Poly-ICLC  
RL: IMF (Industrial manufacture); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

IT 9004-32-4, Carboxymethyl cellulose sodium salt 25104-18-1, Poly-L-lysine 30811-80-4, Poly C 30918-54-8, Polyinosinic acid 38000-06-5, Poly-L-lysine  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(preparation of large volume batches of poly-ICLC with increased biol. potency, and therapeutic use)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 2-9 (Mammalian Hormones)  
Section cross-reference(s): 15

TI Prostaglandin E synthesis and release by murine macrophages and human monocytes after in vitro treatment with biological response modifiers

ST prostaglandin macrophage monocyte biol modifier; lipopolysaccharide prostaglandin macrophage monocyte; polynucleotide complex prostaglandin macrophage monocyte

IT Macrophage  
(PGE formation by, biol. response modifiers effect on)

IT Monocyte  
(PGE formation by, biol. response modifiers effect on human)

IT Lipopolysaccharides  
RL: BIOL (Biological study)  
(PGE formation response to, in macrophage and monocyte of human and laboratory animal)

IT Prostaglandins  
RL: FORM (Formation, nonpreparative)  
(E, formation of, by macrophage and monocyte, of human and laboratory animal,  
biol. response modifiers effect on)

IT Interferons  
( $\alpha$ -, PGE formation response to, in macrophage and monocyte of human and laboratory animal)

IT Interferons  
( $\beta$ -, PGE formation response to, in macrophage and monocyte of human and laboratory animal)

IT 27100-68-1 64118-86-1 64769-70-6 75985-31-8  
RL: BIOL (Biological study)  
(PGE formation response to, in macrophage and monocyte of human and laboratory animal)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 15-5 (Immunochemistry)  
Section cross-reference(s): 1  
TI Production of interferon- $\alpha$  induced by dsRNA in human peripheral  
blood mononuclear cell cultures: role of priming by dsRNA-induced  
interferons- $\gamma$  and - $\beta$   
ST interferon induction doublestranded RNA priming  
IT Ribonucleic acids  
RL: BIOL (Biological study)  
(double-stranded, interferon- $\alpha$  induction by, in human mononuclear  
cells)  
IT Lymphocyte  
Monocyte  
(interaction of, in interferon- $\alpha$  induction)  
IT Leukocyte  
(mononuclear, interferon- $\alpha$  induction in human, interferon- $\beta$   
and - $\gamma$  and cell-cell interactions in)  
IT Interferons  
RL: PRP (Properties)  
( $\alpha$ , induction of, by double-stranded RNA, in human mononuclear  
cells, priming by  $\gamma$ - and  $\beta$ -interferons in)  
IT Interferons  
RL: BIOL (Biological study)  
( $\beta$ , priming by, in  $\alpha$ -interferon induction by double-stranded  
RNA)  
IT Interferons  
RL: BIOL (Biological study)  
( $\gamma$ , priming by, in  $\alpha$ -interferon induction by  
double-stranded RNA)  
IT 24939-03-5, Poly(I):poly(C) 38640-92-5 59789-29-6, Poly(  
ICLC)  
RL: BIOL (Biological study)  
(interferon- $\alpha$  induction by, in human mononuclear cells)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 15-2 (Immunochemistry)  
TI Toll like receptor-3 ligand poly-ICLC promotes the  
efficacy of peripheral vaccinations with tumor antigen-derived peptide  
epitopes in murine CNS tumor models  
ST TLR3 receptor ligand polyICLC CNS tumor vaccine peptide epitope  
IT Proteins  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(EphA2, epitope 671-679; TLR3 receptor ligand poly-  
ICLC promotes efficacy of peripheral vaccination with tumor  
antigen-derived peptide epitopes in murine CNS tumor model)  
IT Toll-like receptors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(TLR-3; TLR3 receptor ligand poly-ICLC promotes  
efficacy of peripheral vaccination with tumor antigen-derived peptide  
epitopes in murine CNS tumor model)  
IT Central nervous system, neoplasm  
Epitopes  
Neuroglia, neoplasm  
(TLR3 receptor ligand poly-ICLC promotes efficacy  
of peripheral vaccination with tumor antigen-derived peptide epitopes  
in murine CNS tumor model)  
IT Peptides, biological studies  
Tumor antigens  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(TLR3 receptor ligand poly-ICLC promotes efficacy  
of peripheral vaccination with tumor antigen-derived peptide epitopes

in murine CNS tumor model)

IT Proteins  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (TRP-2 (tyrosinase-related protein 2), epitope 180-188; TLR3 receptor  
 ligand poly-ICLC promotes efficacy of peripheral  
 vaccination with tumor antigen-derived peptide epitopes in murine CNS  
 tumor model)

IT Peptides, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (comps.; TLR3 receptor ligand poly-ICLC promotes  
 efficacy of peripheral vaccination with tumor antigen-derived peptide  
 epitopes in murine CNS tumor model)

IT T cell (lymphocyte)  
 (cytotoxic; TLR3 receptor ligand poly-ICLC promotes  
 efficacy of peripheral vaccination with tumor antigen-derived peptide  
 epitopes in murine CNS tumor model)

IT Proteins  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (gp100, epitope 25-33; TLR3 receptor ligand poly-ICLC  
 promotes efficacy of peripheral vaccination with tumor antigen-derived  
 peptide epitopes in murine CNS tumor model)

IT Vaccines  
 (tumor; TLR3 receptor ligand poly-ICLC promotes  
 efficacy of peripheral vaccination with tumor antigen-derived peptide  
 epitopes in murine CNS tumor model)

IT Antitumor agents  
 (vaccines; TLR3 receptor ligand poly-ICLC promotes  
 efficacy of peripheral vaccination with tumor antigen-derived peptide  
 epitopes in murine CNS tumor model)

IT Integrins  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 ( $\alpha 4 \beta 1$ ; TLR3 receptor ligand poly-ICLC  
 promotes efficacy of peripheral vaccination with tumor antigen-derived  
 peptide epitopes in murine CNS tumor model)

IT Interferons  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 ( $\gamma$ ; TLR3 receptor ligand poly-ICLC promotes  
 efficacy of peripheral vaccination with tumor antigen-derived peptide  
 epitopes in murine CNS tumor model)

IT 59789-29-6, Poly-ICLC 212370-40-6 219312-69-3  
 841264-18-4  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
 (Biological study); USES (Uses)  
 (TLR3 receptor ligand poly-ICLC promotes efficacy  
 of peripheral vaccination with tumor antigen-derived peptide epitopes  
 in murine CNS tumor model)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 1-7 (Pharmacology)  
 TI Poly ICLC induces anti-IC antibodies in mice and  
 rabbits  
 ST poly ICLC polyinosinate polycytidylate antibody  
 IT Antibodies  
 RL: BIOL (Biological study)  
 (to polyinosinate polycytidylate, poly ICLC  
 induction of)

IT 59789-29-6  
 RL: BIOL (Biological study)  
 (antibodies to polyinosinate polycytidylate production from)

IT 24939-03-5  
 RL: BIOL (Biological study)  
 (antibodies to, poly ICLC induction of)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-6 (Pharmacology)  
TI Drug sensitivity tests against malignant gliomas  
ST antitumor sensitivity test brain glioma  
IT Brain, neoplasm  
(drug sensitivity tests against)  
IT Transplant and Transplantation, animal  
(of glioma, antitumor drug sensitivity test using)  
IT Radiotherapy  
(of gliomas, drug sensitivity tests for)  
IT Neoplasm inhibitors  
(glioma, drug sensitivity tests for)  
IT Neuroglia  
(neoplasm, drug sensitivity tests against)  
IT Lymphokines and Cytokines  
RL: BIOL (Biological study)  
(tumor necrosis factor, glioma inhibition by, drug sensitivity tests for)  
IT Interferons  
RL: BIOL (Biological study)  
( $\alpha$ , glioma inhibition by, drug sensitivity tests for)  
IT Interferons  
RL: BIOL (Biological study)  
( $\beta$ , glioma inhibition by, drug sensitivity tests for)  
IT Interferons  
RL: BIOL (Biological study)  
( $\gamma$ , glioma inhibition by, drug sensitivity tests for)  
IT 50-07-7, Mitomycin C 50-18-0, Cyclophosphamide 57-22-7, Vincristine  
11056-06-7, Bleomycin 15663-27-1, Cisplatin 23214-92-8, Adriamycin  
41598-07-6, Prostaglandin D2 55661-38-6, ACNU 59789-29-6, Poly  
ICLC  
RL: BIOL (Biological study)  
(glioma inhibition by, drug sensitivity tests for)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-5 (Pharmacodynamics)  
Section cross-reference(s): 15  
TI Effect of a nuclease-resistant derivative of polyribonucleosidic-  
polyribocytidylic acid complex on yellow fever in rhesus monkeys (Macaca  
mulatta)  
ST nucleotide complex virucide yellow fever; interferon yellow fever virus  
IT Interferons  
RL: PRP (Properties)  
(induction of, by nucleotide complex, in yellow fever)  
IT Virucides and Virustats  
(nucleotide complex)  
IT Yellow fever  
(nucleotide complex in treatment of)  
IT 59789-29-6  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES  
(Uses)  
(virucidal activity of, in yellow fever)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 15-5 (Immunochemistry)  
TI The in vitro induction of colony-stimulating factor, prostaglandin E, and  
interferon in macrophages and tumor cells by biological response modifiers  
ST colony stimulating factor macrophage tumor cell; prostaglandin E

macrophage tumor cell; interferon macrophage tumor cell; macrophage colony factor prostaglandin interferon; tumor cell colony factor prostaglandin interferon; biol response modifier macrophage tumor cell

IT Neoplasm, metabolism  
(colony-stimulating factor and interferon and prostaglandin E induction in cells of, by biol. response modifiers)

IT Lipopolysaccharides  
RL: BIOL (Biological study)  
(colony-stimulating factor and interferon and prostaglandin E induction in macrophage and tumor cells by)

IT Macrophage  
(colony-stimulating factor and interferon and prostaglandin E induction in, by biol. response modifiers)

IT Interferons  
RL: PRP (Properties)  
(induction of, in macrophage and tumor cells, by biol. response modifiers)

IT Prostaglandins  
RL: PRP (Properties)  
(E, induction of, in macrophage and tumor cells, by biol. response modifiers)

IT 147-84-2, biological studies 27100-68-1 64118-86-1  
RL: BIOL (Biological study)  
(colony-stimulating factor and interferon and prostaglandin E induction in macrophage and tumor cells by)

IT 59789-29-6  
RL: BIOL (Biological study)  
(colony-stimulating factor in and interferon and prostaglandin E induction in macrophage and tumor cells by)

IT 62683-29-8  
RL: PRP (Properties)  
(induction of, in macrophage and tumor cells, by biol. response modifiers)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 1-5 (Pharmacology)  
Section cross-reference(s): 15

TI Characterization of murine Caraparu Bunyavirus liver infection and immunomodulator-mediated antiviral protection

ST hepatitis Caraparu virus treatment immunomodulator; ribavirin interferon hepatitis Caraparu virus

IT Immunostimulants  
(hepatitis from Caraparu virus response to)

IT Virucides and Virustats  
(immunostimulants as, against Caraparu virus-induced hepatitis)

IT Drug interactions  
(of  $\gamma$ -interferons and ribavirin, in hepatitis from Caraparu virus treatment)

IT Virus, animal  
(caraparu, hepatitis from, immunostimulants treatment of)

IT Virus, animal  
(hepatitis, from Caraparu, immunostimulants treatment of)

IT Interferons  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
( $\alpha$ , hepatitis from Caraparu virus response to)

IT Interferons  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
( $\beta$ , hepatitis from Caraparu virus response to)

IT Interferons  
RL: BIOL (Biological study)  
( $\gamma$ , hepatitis from Caraparu virus inhibition by, ribavirin

enhancement of)  
 IT 6307-35-3 27100-68-1 38640-92-5, Ampligen 59789-29-6, Poly  
 ICLC  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); BIOL (Biological study)  
 (hepatitis from Caraparu virus response to)  
 IT 36791-04-5, Ribavirin  
 RL: BIOL (Biological study)  
 (hepatitis from Caraparu virus treatment with,  $\gamma$ -interferon  
 enhancement of)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 1-5 (Pharmacodynamics)  
 Section cross-reference(s): 8  
 TI Evaluation of a nuclease-resistant derivative of poly(I).  
 poly(C) [poly(ICLC)] as a radioprotective  
 agent  
 ST radioprotectant ribonucleic acid complex; polyinosinic polycytidylic  
 complex radioprotectant  
 IT Radioprotectants  
 (polyinosinic acid-polycytidylic acid complex with polylysine and  
 carboxy Me cellulose)  
 IT 59789-29-6  
 RL: BIOL (Biological study)  
 (radioprotectant)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 1-6 (Pharmacology)  
 Section cross-reference(s): 15  
 TI Biological response modifiers: regulators of the cellular immune system  
 and adjuvants in antitumor therapy  
 ST immunomodulator effector cell response antitumor  
 IT Interferons  
 RL: BIOL (Biological study)  
 (cell-mediated immunity response to, and neoplasm inhibition by)  
 IT Immune adjuvants  
 (cell-mediated, neoplasm inhibition by)  
 IT Neoplasm inhibitors  
 (immune adjuvants, cell-mediated immunity in)  
 IT Hematopoiesis  
 Macrophage  
 (neoplasm-inhibiting immune adjuvants effect on)  
 IT Lymphocyte  
 (natural killer, neoplasm-inhibiting immune adjuvants effect on)  
 IT 27100-68-1 39325-01-4 59789-29-6 75985-31-8  
 RL: BIOL (Biological study)  
 (cell-mediated immunity response to, and neoplasm inhibition by)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
 CC 1-5 (Pharmacology)  
 TI Antiviral and immunomodulating inhibitors of experimentally-induced Punta  
 Toro virus infections  
 ST Punto Toro virus virucide immunomodulator  
 IT Immunomodulators  
 Virucides and Virustats  
 (antiviral and immunomodulating inhibitors of exptl.-induced Punta Toro  
 virus infections)  
 IT Polysaccharides, biological studies  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological



study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(glucomannon; antiviral and immunomodulating inhibitors of exptl.-induced Punta Toro virus infections)

IT Virus, animal

(Punta Toro, antiviral and immunomodulating inhibitors of exptl.-induced Punta Toro virus infections)

IT Interferons

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

( $\alpha$ , A/D; antiviral and immunomodulating inhibitors of exptl.-induced Punta Toro virus infections)

IT 54-25-1, 6-Azaauridine 62-53-3, Benzenamine, biological studies

66-81-9, Actidione 145-63-1, Suramin 471-53-4, Glycyrrhetic acid

734-22-5, CL 259763 3930-19-6, Streptonigrin 4016-63-1,

8-Bromoguanosine 6742-12-7, Formycin 12758-40-6, GE132 17073-78-8

19622-83-4, 7-Deoxynarciclasine 25451-90-5 27089-56-1 27100-68-1,

MVE-1 29477-83-6, Narciclasine 29725-42-6 30868-30-5, Pyrazofurin

36703-88-5, Isoprinosine 36791-04-5, Ribavirin 38640-92-5, Ampligen

41729-52-6, 3-Deazaguanine 42400-25-9 56039-11-3, 3-Deazaguanosine

56741-95-8, Bropirimine 58151-87-4 59643-91-3, Imexon 59789-29-6,

Poly(ICLC) 60084-10-8, Tiazofurin 61367-58-6

63166-73-4, Phyllanthoside 68652-43-7, Mannozyim 72161-05-8, Ribavirin

2',3',5'-triacetate 72301-79-2, Enviroxime 81541-26-6, CL 246738

82372-67-6, Pseudolycorine hydrochloride 83161-83-5,

Tiazofurin-5'-monophosphate 83705-13-9, Selenazofurin 87139-86-4, AM 3

87745-28-6, Bryostatin 2 96203-70-2, Pancratistatin 99258-56-7,

Oxamisole 104942-51-0 119567-79-2, Ribamidine 122970-40-5,

7-Thia-8-oxoguanosine 141776-53-6 150316-23-7, Neurotropin

159192-47-9 159192-48-0 159192-49-1

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(antiviral and immunomodulating inhibitors of exptl.-induced Punta Toro virus infections)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN

CC 1-5 (Pharmacodynamics)

Section cross-reference(s): 15

TI Modified polyribonucleosinic-polyribocytidylic acid complex: induction of serum interferon, fever, and hypotension in rabbits

ST polyinosinate polycytidylate polylysine complex pharmacol; interferon

polyinosinate polycytidylate polylysine; fever polyinosinate

polycytidylate polylysine; hypotension polyinosinate polycytidylate

polylysine

IT Fever and Hyperthermia

Hypotension

(from poly(I)-poly(C)-poly(L-lysine) complex, interferon induction in relation to)

IT Interferons

RL: BIOL (Biological study)

(induction by, poly(I)-poly(C)-poly(L-lysine) complex, fever and hypotension in relation to)

IT 24939-03-5D, poly-L-lysine complex 25104-18-1D, poly(I)-poly(C) complex

38000-06-5D, poly(I)-poly(C) complex

RL: BIOL (Biological study)

(fever and hypotension and interferon induction by)

IT 50-23-7

RL: BIOL (Biological study)

(fever and hypotension from poly(I)-poly(C)-poly(L-lysine) complex decrease by, interferon induction in relation to)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L1 94 ANSWERS CAPLUS COPYRIGHT 2007 ACS on STN  
CC 1-5 (Pharmacology)  
TI Enhanced therapeutic efficacy of poly(ICLC) and  
ribavirin combinations against Rift Valley fever virus infection in mice  
ST virus infection polyICLC ribavirin  
IT Virus, animal  
(Rift Valley fever, infection with, therapeutic efficacy of  
poly(ICLC) and ribavirin combinations against)  
IT 36791-04-5, Ribavirin  
RL: BIOL (Biological study)  
(Rift Valley fever virus infection response to poly(  
ICLC) and)  
IT 59789-29-6, Poly ICLC  
RL: BIOL (Biological study)  
(Rift Valley fever virus infection treatment with ribavirin and)

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> d his

(FILE 'HOME' ENTERED AT 18:43:07 ON 11 JUL 2007)

FILE 'CAPLUS' ENTERED AT 18:43:17 ON 11 JUL 2007

L1 94 S (POLY (5A) ICLC)  
L2 2 L1 AND PREP/RL

=> s l1 and lysine and (carboxymethylcellulose OR "Carboxymethyl cellulose")

108942 LYSINE  
2345 LYSINES  
109676 LYSINE  
(LYSINE OR LYSINES)  
7316 CARBOXYMETHYLCELLULOSE  
62 CARBOXYMETHYLCELLULOSES  
7344 CARBOXYMETHYLCELLULOSE  
(CARBOXYMETHYLCELLULOSE OR CARBOXYMETHYLCELLULOSES)  
37332 "CARBOXYMETHYL"  
3 "CARBOXYMETHYLS"  
37332 "CARBOXYMETHYL"  
( "CARBOXYMETHYL" OR "CARBOXYMETHYLS")  
355188 "CELLULOSE"  
4389 "CELLULOSES"  
355685 "CELLULOSE"  
( "CELLULOSE" OR "CELLULOSES")  
13275 "CARBOXYMETHYL CELLULOSE"  
( "CARBOXYMETHYL" (W) "CELLULOSE")

L3 25 L1 AND LYSINE AND (CARBOXYMETHYLCELLULOSE OR "CARBOXYMETHYL  
CELLULOSE")

=> d l3 1-25 ibib abs

L3 ANSWER 1 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:18891 CAPLUS

DOCUMENT NUMBER: 140:71067

TITLE: Method for preparation of large volume batches of  
poly-ICLC with increased biological  
potency, and therapeutic, clinical and veterinary uses  
thereof

INVENTOR(S): Salazar, Andres

PATENT ASSIGNEE(S): Oncovir, Inc., USA

SOURCE: U.S. Pat. Appl. Publ., 14 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2004005998	A1	20040108	US 2003-611614	20030701
WO 2005102278	A1	20051103	WO 2003-US20828	20030701
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2003248791	A1	20051109	AU 2003-248791	20030701
EP 1778186	A1	20070502	EP 2003-819324	20030701
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PT, RO, SE, SI, SK, TR				

PRIORITY APPLN. INFO.: US 2002-393713P P 20020703  
WO 2003-US20828 W 20030701

AB The invention discloses a method for producing large lots of final sterile poly-ICLC suitable for clin. use with reduced toxicity at ED levels, as well as a method for using poly-ICLC to regulate genes and a method for using poly-ICLC to treat certain human and veterinary infectious, neoplastic and autoimmune disorders.

L3 ANSWER 2 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:216000 CAPLUS

DOCUMENT NUMBER: 108:216000

TITLE: Interferon induction in piglets with polyinosinic:polycytidylic acid complexed with poly-L-lysine and carboxymethylcellulose

AUTHOR(S): Loewen, K. G.; Derbyshire, J. B.

CORPORATE SOURCE: Dep. of Vet., Univ. Guelph, Guelph, N1G 2W1, Can.

SOURCE: Research in Veterinary Science (1988), 44(1), 132-3

CODEN: RVTSA9; ISSN: 0034-5288

DOCUMENT TYPE: Journal

LANGUAGE: English

AB When newborn piglets were inoculated i.v. with 1.0, 0.5 or 0.25 mg kg<sup>-1</sup> of poly I:C complexed with poly-L-lysine and CM-cellulose (poly ICLC), the highest serum interferon levels and the lowest white blood cell counts were found in response to a dose of 0.5 mg kg<sup>-1</sup>. Similar responses were observed in weaned piglets inoculated with 0.25 mg kg<sup>-1</sup> of poly ICLC. Poly ICLC was a more effective interferon inducer than poly I:C, particularly in newborn piglets.

L3 ANSWER 3 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:142978 CAPLUS

DOCUMENT NUMBER: 108:142978

TITLE: Increase in liver-associated natural killer activity by polyribonucleotides

AUTHOR(S): Twilley, Theresa A.; Mason, Lewellyn; Talmadge, James E.; Wilttrout, Robert H.

CORPORATE SOURCE: Lab. Exp. Immunol., Biol. Response Modifiers Program, Frederick, MD, USA

SOURCE: Natural Immunity and Cell Growth Regulation (1987), 6(6), 279-90

CODEN: NICRDR; ISSN: 0254-7600

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In mice, polyinosinic-polycytidylic acid and poly-L-lysine which has been stabilized in carboxymethylcellulose (polyICLC) as well as polyadenosinic-polyuridylic acid (poly AU), both potently augmented natural killer (NK) activity in the liver. Following the administration of poly ICLC (10 µg/mouse) greater NK activity, as measured by lytic units (LU), was observed in the liver (445 LU) than in blood (63 LU) or spleen (20 LU). The high level of NK activity in the liver was in contrast to the low levels observed in untreated mice, and was maintained for at least 9 days post injection. NK activity in the blood and spleen returned to normal levels by day 6. Similar results were obtained with polyAU except that approx. 10-fold more poly AU (100 µg/mouse) was required to induce optimal augmentation of NK activity. The increase in liver-associated NK activity induced by poly ICLC was associated with a 10- to 20-fold increase in liver-associated leukocytes, termed nonparenchymal cells (NPC). The NK activity mediated by NPC was associated with cells morphol. characterized as large granular lymphocytes (LGL). The repeated administration of poly ICLC resulted in higher levels of liver-associated NK activity and total liver-associated LGL as compared to a single injection.

L3 ANSWER 4 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:15951 CAPLUS

DOCUMENT NUMBER: 108:15951

TITLE: Toxicity of polyinosinic-polycytidylic acid admixed with poly-L-lysine and solubilized with carboxymethylcellulose in mice

AUTHOR(S): Hartmann, Diethelm; Schneider, Mark A.; Lenz, Barbara F.; Talmadge, James E.

CORPORATE SOURCE: Preclin. Screening Lab., Natl. Cancer Inst., Frederick, MD, USA

SOURCE: Pathology and Immunopathology Research (1987), 6(1), 37-50

CODEN: PIREEI; ISSN: 0257-2761

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Polyinosinic-polycytidylic acid admixed with poly-L-lysine and solubilized with carboxymethylcellulose (poly(I,C)-LC) reduced the body wts. of treated mice and induced hepatic necrosis and pulmonary toxicity after i.p. and i.v. administrations, resp. The body wts. recovered with time despite repeated treatments.

L3 ANSWER 5 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:205212 CAPLUS

DOCUMENT NUMBER: 104:205212

TITLE: Potentiation of the cytotoxic effect of human immune interferon by different synthetic double-stranded RNAs in the refractory human colon carcinoma cell line BE

AUTHOR(S): Chapekar, Mrunal S.; Glazer, Robert I.

CORPORATE SOURCE: Lab. Biol. Chem., Natl. Cancer Inst., Bethesda, MD, 20892, USA

SOURCE: Cancer Research (1986), 46(4, Pt. 1), 1698-702

CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A human cell line BE, derived from an undifferentiated carcinoma of the colon, was studied for its response to the cytotoxic effects of human immune interferon (IFN-γ) alone and in combination with various double-stranded RNAs (dsRNAs). BE cells were moderately refractory to 3-day treatment with IFN-γ (10-300 units/mL) where only 5-30% reduction in colony formation occurred. A similar exposure interval to poly(I)·poly(C) (100 µg/mL) had no detectable effect on colony formation. In contrast, the lethal effect of the combination of IFN-γ and poly(I)·poly(C) was synergistic and this regimen produced a 40-80% reduction in colony formation. The cytotoxic effects of the combination of IFN-γ with varying concns. of the dsRNAs

poly(I)·poly(C), poly(A)·poly(U), polyinosinic·polyribocytidylic acid stabilized with poly-L-lysine in carboxymethylcellulose [poly(ICLC)], and mismatched dsRNA [rIn·r(C13,U)n] were also examined. The concentration of the dsRNAs producing a 50% decrease in cell viability in combination with IFN-γ (100 units/mL) was 6 μg/mL for poly(I)·poly(C), 1 μg/mL for poly(A)·poly(U), 3 ng/mL for poly(ICLC), and 16 μg/mL for rIn·r(C13,U)n. DNA, RNA, and protein synthesis in IFN-γ and poly(I)·poly(C)-treated cells were reduced in a dose-dependent manner. There were no changes in either (2',5')oligoadenylate concns. or in rRNA transcription following treatment with IFN-γ and poly(I)·poly(C). Thus, the synergism resulting from the combination of IFN-γ and dsRNA appears to be mediated via another, as yet unknown, mechanism.

L3 ANSWER 6 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:14714 CAPLUS  
DOCUMENT NUMBER: 104:14714  
TITLE: Purification of carboxymethylcellulose decreases toxicity of poly ICLC in mice

AUTHOR(S): Bello, Jake; O'Malley, Judith; Granados, Edward  
CORPORATE SOURCE: Dep. Biophys., Roswell Park Mem. Inst., Buffalo, NY, 14263, USA

SOURCE: Journal of Interferon Research (1985), 5(3), 429-30  
CODEN: JIREDJ; ISSN: 0197-8357

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Purification of the carboxymethylcellulose (CMC) component of polyinosinic polycytidylic acid · poly-L-lysine · carboxymethylcellulose (I) [59789-29-6] by EtOH extraction did not significantly affect the efficacy of I as an interferon inducer in mice, but decreased the toxic side effects.

L3 ANSWER 7 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:14713 CAPLUS  
DOCUMENT NUMBER: 104:14713  
TITLE: A comparison of interferon responses to poly ICLC in males and females

AUTHOR(S): Bever, Christopher T., Jr.; McFarlin, Dale E.; Levy, Hilton B.

CORPORATE SOURCE: Neuroimmunol. Branch, NINCDS, Bethesda, MD, USA

SOURCE: Journal of Interferon Research (1985), 5(3), 423-8  
CODEN: JIREDJ; ISSN: 0197-8357

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Interferon (IFN) responses to polyriboinosinic acid polyribocytidylic acid · poly-L-lysine · carboxymethylcellulose (poly ICLC) [59789-29-6] were studied in humans as part of a preliminary trial in patients with multiple sclerosis (MS). Patients received i.v. doses of 100 μg/kg poly ICLC. Men and women produced substantial levels of IFN at 8, 12, and 16 h after infusion, but the levels of IFN in men were consistently higher. Interferon responses were also examined in male and female Rhesus monkeys. Again, there were higher levels of IFN in males. The observed differences may reflect sex-linked differences in either drug metabolism or specific sensitivity to IFN induction by poly ICLC. The most interesting possibility is that the difference is due to a more general difference in IFN response between males and females.

L3 ANSWER 8 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1985:605785 CAPLUS  
DOCUMENT NUMBER: 103:205785

TITLE: Response of mouse tumor to interferon inducer and radiation  
AUTHOR(S): Lvovsky, Edward A.; Mossman, Kenneth L.; Levy, Hilton B.; Dritschilo, Anatoly  
CORPORATE SOURCE: Med. Cent., Georgetown Univ., Washington, DC, 20007, USA  
SOURCE: International Journal of Radiation Oncology, Biology, Physics (1985), 11(9), 1721-5  
CODEN: IOBPD3; ISSN: 0360-3016  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The antitumor effect of interferon inducer poly(ICLC) (polyriboinosinic acid-polyribocytidylic acid-poly-L-lysine-carboxymethylcellulose complex) [59789-29-6], given prior to radiation treatment of Lewis lung carcinoma in C57B1 mice was studied. The local response, as measured by the delay in the tumor growth, was significantly higher in the combination treatment group than in poly(ICLC) or local irradiation groups. Following the termination of treatment, tumor regrowth was observed. The survival of poly(ICLC) treated mice was influenced by the number of transplanted tumor cells. Thus, untreated mice which received  $3 + 104$  or  $3 + 105$  (2 or 20 TD50) of tumor cells had similar mean survival time of 25.4 and 22 days, resp. The mice, treated by a combination of poly(ICLC) and local irradiation survived 48.2 days and 30.7 days, with higher survival in 2 TD50 tumor cell groups. Thus, data obtained in this study in mice showed that administration of an interferon inducer polyn(ICLC) prior to local irradiation can improve survival.

L3 ANSWER 9 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1985:469574 CAPLUS  
DOCUMENT NUMBER: 103:69574  
TITLE: The in vitro induction of colony-stimulating factor, prostaglandin E, and interferon in macrophages and tumor cells by biological response modifiers  
AUTHOR(S): Schlick, Erich; Hartung, Klaus; Piccoli, Mario; Bartocci, Anna; Chirigos, Michael A.  
CORPORATE SOURCE: Frederick Cancer Res. Facil., Natl. Cancer Inst., Frederick, MD, USA  
SOURCE: Immunology Series (1984), 25(Immune Modulation Agents Their Mech.), 513-29  
CODEN: IMSED7; ISSN: 0092-6019  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The effects of 6 biol. response modifiers (BRMs) on the secretion of colony-stimulating factors (CSFs), prostaglandin E, and interferon were studied using resident peritoneal macrophages (M0) and 2 tumor cell lines, Wehi-3 and L1210, in an in vitro system. The BRMs most effective for M0 were lipopolysaccharide (LPS), interferon (IF), and polyriboinosinic acid-polycytidylic acid poly-L-lysine stabilized with carboxymethylcellulose (poly ICLC). However, only poly ICLC induced considerable amts. of IF in M0. The same 3 BRMs induced CFS production by the 2 tumor cell lines whereas none of the drugs could induce prostaglandin E and only poly ICLC stimulated marginal IF titers (.apprx.10 units/mL) in the cell lines.

L3 ANSWER 10 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1985:464451 CAPLUS  
DOCUMENT NUMBER: 103:64451  
TITLE: Cellular regulation by immunomodifiers MVE-2 and poly ICLC and their therapeutic application  
AUTHOR(S): Chirigos, Michael A.; Saito, Tohru; Schlick, Erich; Ruffman, Ralf

CORPORATE SOURCE: Frederick Cancer Res. Fac., Natl. Cancer Inst.,  
Frederick, MD, 21701, USA  
SOURCE: NIH Publ. (1985), n85-1177, Cancer Treat. Symp., 1985,  
Vol. 1, 11-18  
CODEN: DPNSDO  
DOCUMENT TYPE: Report  
LANGUAGE: English

AB Two immunomodifiers, maleic anhydride divinyl ether copolymer (MVE-2) [27100-68-1] and polyinosinic-polycytidylic acid poly-L-lysine stabilized with carboxymethylcellulose (poly ICLC) [59789-29-6], augmented natural killer (NK) cell activity in several tissues. Macrophage tumoricidal activity was also markedly increased. Both effector cells were active for 1 wk, with macrophage activity remaining elevated for a longer period. Multiple treatment with both agents resulted in a decrease in NK cell response, but macrophage activity remained elevated; NK cells had the greatest hyporesponsiveness to MVE-2. Both agents caused an increase in secretion of colony-stimulating factor from bone marrow cells and in serum. Treatment with MVE-2 and poly ICLC resulted in an earlier reconstitution of bone marrow cells, NK cell activity, and macrophage effector cell activity in mice pretreated with cyclophosphamide [50-18-0]; MVE-2 prevented the establishment of B16 melanoma metastasis in lung and liver. Thus, NK cells and macrophages play a supportive role in the natural resistance to tumor cell replication. Combined treatment of MBL-2 tumor cells with cytoreductive chemotherapy with cyclophosphamide and MVE-2 resulted in an enhanced therapeutic response.

L3 ANSWER 11 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1985:432069 CAPLUS  
DOCUMENT NUMBER: 103:32069  
TITLE: Poly ICL-CM dextran: an interferon inducer of reduced toxicity  
AUTHOR(S): Granados, Edward N.; Dawidzik, Jean; O'Malley, Judith; McGarry, Michael; Bello, Jake  
CORPORATE SOURCE: Abbott Lab., North Chicago, IL, 60064, USA  
SOURCE: Journal of Interferon Research (1984), 4(2), 155-60  
CODEN: JIREDJ; ISSN: 0197-8357  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Poly ICL-CM dextran [97127-12-3] (a complex of polyinosinate-polycytidylate, poly-L-lysine, and carboxymethyl dextran) was as effective an inducer of interferon in mice and rhesus monkeys as poly ICLC (the counterpart of poly ICL-CM dextran which contains carboxymethylcellulose instead of carboxymethyl dextran). In addition, poly ICL-CM dextran showed lower toxicity in mice than poly ICLC.

L3 ANSWER 12 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1983:137376 CAPLUS  
DOCUMENT NUMBER: 98:137376  
TITLE: Inhibition of virus-induced murine diabetes by an interferon inducer  
AUTHOR(S): Gadzik, James P.; Naji, Ali; Barker, Clyde F.; Blank, Kenneth J.  
CORPORATE SOURCE: Sch. Med., Univ. Pennsylvania, PA, USA  
SOURCE: Journal of Interferon Research (1982), 2(1), 59-63  
CODEN: JIREDJ; ISSN: 0197-8357  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Administration of the interferon inducer poly (I)·poly(C)-poly-L-lysine in carboxymethylcellulose (I) prior to inoculation with the picornavirus, encephalomyocarditis virus (EMCV), and at 48 and 96 h thereafter effectively blocked the induction of diabetes in mice during a 36-day period. Pretreatment with a single dose of I prior to virus inoculation afforded protection during the 1st wk after infection

(as indicated by a decreased hyperglycemia), but was a transient effect.

L3 ANSWER 13 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1982:504227 CAPLUS  
DOCUMENT NUMBER: 97:104227  
TITLE: Poly(ICL) as an effective interferon inducer  
INVENTOR(S): Levy, Hilton B.; Riley, Freddie L.  
PATENT ASSIGNEE(S): United States Dept. of Health and Human Services, USA  
SOURCE: U. S. Pat. Appl., 24 pp. Avail. NTIS Order No.  
PAT-APPL-6-292 583.  
CODEN: XAXXAV  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 292583	A0	19820618	US 1981-292583	19810831
PRIORITY APPLN. INFO.:			US 1981-292583	19810831

AB A polyribonucleosinic-polyribocytidylic-poly-L-lysine complex [poly(ICL)] [35560-71-5] containing different amts. of poly-L-lysine [poly(L)] was prepared by dropwise addition at 58-60° of different vols. of a poly(L) solution to a solution which had been prepared from equal amts. of polyribonucleosinic [poly(I)] and polyribocytidylic [poly(C)] stock solns. The poly(ICL) complexes were as effective interferon inducers in mice as was a previously described poly(I:C)-poly(L)-carboxymethylcellulose [poly(ICLC)] complex, and more effective than poly(I:C). Poly(ICL) was less toxic in mice than was poly(ICLC) (i.p. LD50 values of 25.1 and 12.6 mg/kg, resp.). The poly(ICL) complex with the highest amount of poly(L) induced the highest levels of interferon in Rhesus monkeys. Advantages of the poly(ICL) complex over the poly(ICLC) complex are discussed.

L3 ANSWER 14 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1982:503984 CAPLUS  
DOCUMENT NUMBER: 97:103984  
TITLE: Interferon induction and therapy of brain tumors in rats by poly(ICLC)  
AUTHOR(S): Machida, Haruhiko; Takezawa, Junichi; Kuninaka, Akira; Yoshino, Hiroshi; Nakamura, Osamu; Takakura, Kintomo  
CORPORATE SOURCE: Res. Lab., Yamasa Shoyu Co. Ltd., Choshi, 288, Japan  
SOURCE: Microbiology and Immunology (1982), 26(4), 353-8  
CODEN: MIIMDV; ISSN: 0385-5600  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The levels of plasma interferon in rats 4 h after injection of several doses (0.05-5 mg/kg) of poly(ICLC)-B or -C [poly(I)·poly(C) stabilized with varying amts. of poly-L-lysine-carboxymethylcellulose [59789-29-6]] were the same as those in rats receiving equal doses of poly(I)·poly(C) [24939-03-5]. (The amount of inducer is expressed in terms of poly(I)·poly(C) content of each preparation). However, the interferon level of plasma persisted for a longer time in rats injected with poly(ICLC)-B than in those injected with just poly(I)·poly(C). Treatment with poly(ICLC)-B (i.v.) was moderately effective in increasing the survival time of rats inoculated intracerebrally with glial tumor cells when the treatment was started by 7 days after tumor cell inoculation. Thus, the antitumor activity of poly(ICLC)-B is correlated with persistence of the high level of interferon induced thereby, and, probably also with the immune adjuvant activity of poly(ICLC).

L3 ANSWER 15 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN



ACCESSION NUMBER: 1982:135517 CAPLUS  
DOCUMENT NUMBER: 96:135517  
TITLE: Polyribonucleosinic-polyribocytidylic acid-poly-L-lysine complex [poly(ICL)] without carboxymethylcellulose (CMC): a new primate-effective interferon inducer  
AUTHOR(S): Riley, Freddie L.; Morin, Martin L.; Lvovsky, Eduard; Stephens, Edward E.; Levy, Hilton B.  
CORPORATE SOURCE: Natl. Inst. Allergy Infect. Dis., Bethesda, MD, 21701, USA  
SOURCE: Proceedings of the Society for Experimental Biology and Medicine (1982), 169(2), 183-8  
CODEN: PSEBAA; ISSN: 0037-9727  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The complex of poly(I).poly(C) with poly-(L-lysine) in 0.5% carboxymethylcellulose (CMC) [poly(ICLC)] has proven to be an effective interferon inducer in primates, including man. Since no mechanism is known by which the body can degrade CMC, a new complex of lower mol. weight, which contains poly I.poly C complexed with poly-L-lysine [poly(ICL)], but without CMC, was developed. This compound is slightly more resistant than poly(ICLC) to hydrolysis by RNase A and is also an effective inducer of interferon in nonhuman primates. The new compound without CMC is also less toxic in mice than is poly(ICLC) as indicated by LD50 values.

L3 ANSWER 16 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1982:83980 CAPLUS  
DOCUMENT NUMBER: 96:83980  
TITLE: Immune response modifying activity in mice of polyinosinic: polycytidylic acid stabilized with poly-L-lysine, in carboxymethylcellulose [poly-ICLC]  
AUTHOR(S): Chirigos, M. A.; Papademetriou, V.; Bartocci, A.; Read, E.; Levy, H. B.  
CORPORATE SOURCE: Div. Cancer Treat., Natl. Cancer Inst., Bethesda, MD, 20205, USA  
SOURCE: International Journal of Immunopharmacology (1981), 3(4), 329-37  
CODEN: IJIMDS; ISSN: 0192-0561  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Poly-ICLC, a polyinosinic polycytidylic acid stabilized with poly-L-lysine in carboxymethylcellulose, was tested in mice for its immunoregulatory activity. Poly-ICLC enhanced T cell responsiveness but not B cell responsiveness. It augmented the delayed type hypersensitivity response significantly. The results indicate that Poly-ICLC is a T cell stimulator. Macrophage tumoricidal activity was markedly enhanced both in vitro and in vivo after exposure to Poly-ICLC. Natural killer cell cytotoxicity was significantly augmented in vivo. Both macrophage and natural killer cell activity was maintained for over 3 days after only one treatment. The extended period of tumor cell cytotoxicity, exhibited by macrophage and natural killer cells, may correlate to Poly-ICLC induction of early and high levels of interferon which are maintained in the serum for a longer period of time.

L3 ANSWER 17 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1980:597823 CAPLUS  
DOCUMENT NUMBER: 93:197823  
TITLE: Modified polyribonucleosinic-polyribocytidylic acid complex: modulation of toxicity for rabbits by alterations in components

AUTHOR(S): Gatmaitan, Bienvenido G.; Levy, Hilton B.; Lerner, A. Martin  
CORPORATE SOURCE: VA Med. Cent., Allen Park, MI, 48101, USA  
SOURCE: Antimicrobial Agents and Chemotherapy (1980), 18(3), 409-15  
CODEN: AMACCQ; ISSN: 0066-4804  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The effects of a modified complex of polyriboinosinic acid-polyribocytidylic acid [poly(I)-poly(C)] with carboxymethylcellulose and poly-L-lysine [poly(ICLC) [59789-29-6]], namely, the induction of high titers of serum interferon along with fever and hypotension, were reproduced in rabbits. The effects of complexes with homopolymer polyribonucleotide sedimentation coeffs. decreasing from 9S to 6S and 4S were evaluated. Modified complexes of carboxymethylcellulose and poly-L-lysine with decreasing mol. wts. were also tested. In several studies diphenhydramine [58-73-1] and indomethacin [53-86-1] were administered concomitantly. At a daily i.v. dose of 0.2 mg/kg, the various preps. of poly(I)-poly-(C) (9S) and poly(I)-poly(C) (6S) induced falls in blood pressure, but stabilized complexes of poly(I)-poly(C) (4S) did not. Alterations in the mol. wts. of carboxymethylcellulose and poly-L-lysine in the modified complex and the concomitant administration of diphenhydramine did not influence the occurrence or severity of untoward reactions. In rabbits, poly(ICLC) (4S) along with indomethacin induced high titers of serum interferon without fever or hypotension.

L3 ANSWER 18 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1980:507229 CAPLUS

DOCUMENT NUMBER: 93:107229

TITLE: Interferon induction by and toxicity of polyriboinosinic acid [poly(rI)].polyribocytidylic acid [poly(rC)], mismatched analog poly(rI).poly[r(C12uracil)n], and poly(rI).poly(rC) L-lysine complexed with carboxymethylcellulose

AUTHOR(S): Stringfellow, Dale A.; Weed, Sheldon D.  
CORPORATE SOURCE: Exp. Biol. Res., Upjohn Co., Kalamazoo, MI, 49001, USA  
SOURCE: Antimicrobial Agents and Chemotherapy (1980), 17(6), 988-92  
CODEN: AMACCQ; ISSN: 0066-4804

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The ability of polyriboinosinic acid-polyribocytidylic acid (poly I-polyC) [24939-03-5], mismatched analog poly I-poly[(C12Uracil)n] [38640-92-5], and poly I-poly C complexed with poly L-lysine and carboxymethylcellulose [poly(ICLC)] [59789-29-6] to induce interferon and the comparative toxicity of each in cats were evaluated. Each induced high levels of circulating interferon, although poly(ICLC) injected i.v. at 1 to 4 mg/kg induced  $\leq 10$ -fold more interferon than the other compds. Each compound was pyrogenic and caused a transient decrease in leukocyte nos. Poly I-polyC and the mismatched analog caused severe diarrhea and nausea at the highest drug concns. (1 to 4 mg/kg), but poly(ICLC) did not. Each compound also caused depression and lethargy and impaired coordination.

L3 ANSWER 19 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1979:418239 CAPLUS

DOCUMENT NUMBER: 91:18239

TITLE: Adjuvant effects of low doses of a nuclease-resistant derivative of polyinosinic acid-polycytidylic acid on antibody responses of monkeys to inactivated

AUTHOR(S): Venezuelan equine encephalomyelitis virus vaccine  
Harrington, D. G.; Crabbs, C. L.; Hilmas, D. E.;  
Brown, J. R.; Higbee, G. A.; Cole, F. E., Jr.; Levy,  
H. B.  
CORPORATE SOURCE: Army Med. Res. Inst. Infect. Dis., Fort Detrick,  
Frederick, MD, 21701, USA  
SOURCE: Infection and Immunity (1979), 24(1), 160-6  
CODEN: INFIBR; ISSN: 0019-9567  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Polyriboinosinic-polyribocytidylic acid [poly(I)·poly(C)]  
stabilized with poly-L-lysine and CM-cellulose [poly(  
ICLC)] has been previously shown to be a compound with marked  
adjuvant activity when given in high doses with inactivated Venezuelan  
equine encephalomyelitis (VEE) virus vaccine. This study investigated the  
effects of much lower doses of poly(ICLC) on the  
magnitude and kinetics of the primary and secondary humoral antibody  
responses of rhesus monkeys to inactivated VEE virus vaccine. Monkeys  
given a single injection of vaccine developed very low neutralizing  
antibody titers, whereas those given adjuvant plus vaccine had  
30-100-fold-higher titers which remained elevated for >6 mo. Low doses of  
poly(ICLC) given with VEE virus vaccine resulted in a  
profound but transient increase in priming of secondary antibody responses  
to the antigen. In contrast, the administration of poly-L-lysine  
and CM-cellulose alone without the poly(I)·poly(C) component of the  
complex had no adjuvant effect on antibody responses of monkeys to VEE  
virus vaccine. The temporal development of antibody by class (IgM-IgG) in  
monkeys given 2 injections of adjuvant-vaccine was not different from that  
with vaccine alone. Serial hematol. and clin. chemical detns. on monkeys  
given single or multiple doses of poly(ICLC) with  
vaccine were not different from values in monkeys given vaccine alone.

L3 ANSWER 20 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1979:400321 CAPLUS  
DOCUMENT NUMBER: 91:321  
TITLE: Protective and toxic effects of a nuclease-resistant  
derivative of polyriboinosinic-polyribocytidylic acid  
on Venezuelan equine encephalomyelitis virus in Rhesus  
monkeys  
AUTHOR(S): Stephen, E. L.; Hilmas, D. E.; Levy, H. B.; Spertzel,  
R. O.  
CORPORATE SOURCE: Army Med. Res. Inst. Infect. Dis., NIH, Bethesda, MD,  
USA  
SOURCE: Journal of Infectious Diseases (1979), 139(3), 267-72  
CODEN: JIDIAQ; ISSN: 0022-1899  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Poly I-poly C [24939-03-5], stabilized with poly-L-  
lysine and carboxymethylcellulose (poly  
ICLC), favorably altered the pathogenesis of Venezuelan equine  
encephalomyelitis virus infection in rhesus monkeys by decreasing the number  
of infected monkeys that became detectably viremic and by delaying the  
onset of viremia in the remaining monkeys. The death of some infected,  
treated monkeys in the absence of death in monkeys that were either  
infected and untreated or treated and uninfected suggests a synergistic  
toxicity resulting from the combination of infection, handling, and  
poly ICLC treatment, although other explanations are  
possible.

L3 ANSWER 21 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1978:523224 CAPLUS  
DOCUMENT NUMBER: 89:123224  
TITLE: Effect of interferon on togavirus and arenavirus  
infections of animals  
AUTHOR(S): Stephen, E. L.; Scott, S. K.; Eddy, G. A.; Levy, H. B.

CORPORATE SOURCE: U. S. Army Med. Res. Inst. Infect. Dis., Fort Detrick,  
MD, USA  
SOURCE: Texas Reports on Biology and Medicine (1977),  
35(Interferon Syst.), 449-54  
CODEN: TRBMAV; ISSN: 0040-4675  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Machupo virus-infected monkeys treated early in infection with the  
interferon inducer poly(ICLC) [poly  
(I).cntdot.poly(C)-carboxymethylcellulose-poly-L-  
lysine complex] had significantly higher viremias than did  
untreated, control monkeys. A possible explanation for the higher  
viremias is that interferon and/or poly(ICLC) either  
altered or stimulated the production of certain cell types that are target  
tissues for viral replication. The results are discussed with respect to  
the sensitivity of togavaruses and arenaviruses to interferon.

L3 ANSWER 22 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1978:83623 CAPLUS  
DOCUMENT NUMBER: 88:83623  
TITLE: Use of poly(ICLC) for the  
prophylaxis and treatment of Venezuelan equine  
encephalomyelitis virus infection in nonhuman primates  
AUTHOR(S): Hilmas, Duane E.; Stephen, Edward L.; Spertzel,  
Richard O.; Levy, Hilton B.  
CORPORATE SOURCE: Army Med. Res. Inst. Infect. Dis., Frederick, MD, USA  
SOURCE: U. S. NTIS, AD Rep. (1977), AD-A044417, 13 pp.  
Avail.: NTIS  
From: Gov. Rep. Announce. Index (U. S.) 1977, 77(24),  
92  
CODEN: XADRCH; ISSN: 0099-8575  
DOCUMENT TYPE: Report  
LANGUAGE: English

AB Poly(I).poly(C) stabilized with poly-L-lysine  
and carboxymethylcellulose [poly(ICLC)]  
[59789-29-6] induced moderate to high levels of serum interferon in man,  
nonhuman primates, and rodents. Poly(ICLC) was tested  
therapeutically in rhesus monkeys against infection with a virulent strain  
of Venezuelan equine encephalomyelitis (VEE) virus. The VEE-1 strain of  
virus in exptl. infections presents a broad spectrum of interactions with  
different hosts, ranging from mild, subclin. infections to severe,  
prostrating disease and death. In monkeys, clin. signs of exptl.  
infection are generally of a mild form, with a characteristic biphasic  
febrile response clearly detectable but nonlethal. This strain of virus  
is sensitive to interferon in vitro. Several monkeys inoculated with 1000  
plaque-forming units or more of VEE-1 virus died (11-20 days) subsequent  
to infection when treated with 3.0 mg/kg of poly(ICLC  
) . Poly(ICLC) is reported to be an effective  
antiviral agent against lethal yellow fever and Japanese encephalitis  
virus infections in monkeys. The variable response found following  
treatment of VEE virus disease in monkeys suggests caution regarding the  
indiscriminate use of poly(ICLC) to treat virus  
infections that have not been specifically evaluated with respect to their  
response to this drug.

L3 ANSWER 23 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN  
ACCESSION NUMBER: 1978:292 CAPLUS  
DOCUMENT NUMBER: 88:292  
TITLE: Effect of a nuclease-resistant derivative of  
polyribonucleosinic-polyribocytidylic acid complex on  
yellow fever in rhesus monkeys (Macaca mulatta)  
AUTHOR(S): Stephen, E. L.; Sammons, M. L.; Pannier, W. L.; Baron,  
S.; Spertzel, R. O.; Levy, H. B.  
CORPORATE SOURCE: U. S. Army Med. Res. Inst. Infect. Dis., Fort Detrick,  
Frederick, MD, USA

SOURCE: Journal of Infectious Diseases (1977), 136(1), 122-6  
CODEN: JIDIAQ; ISSN: 0022-1899

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Rhesus monkeys (*Macaca mulatta*) treated with a newly developed nuclease-resistant polyribonucleosinic-polyribocytidylic acid-poly-L-lysine-carboxymethylcellulose complex [poly (ICLC)] [64685-78-5] did not die after challenge with virulent Asibi strain yellow fever (YF) virus. The strain of virus is sensitive to the effects of interferon in vitro and is lethal for rhesus monkeys 4 to 6 days after s.c. administration of 1,000 plaque-forming units of the virus. The mortality rate was reduced in monkeys initially treated 8 h before or after inoculation of virus but was unchanged in monkeys initially treated 24 h after challenge. Treated monkeys developed neutralizing antibody to YF virus. The successful treatment of yellow fever in a primate model with use of poly (ICLC) suggests a meaningful role for the interferon system in the host defense against this viral infection.

L3 ANSWER 24 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1977:565851 CAPLUS

DOCUMENT NUMBER: 87:165851

TITLE: Swine influenza virus vaccine: potentiation of antibody responses in rhesus monkeys

AUTHOR(S): Stephen, E. L.; Hilmas, D. E.; Mangiafico, J. A.; Levy, H. B.

CORPORATE SOURCE: U. S. Army Med. Res. Inst. Infect. Dis., Frederick, MD, USA

SOURCE: Science (Washington, DC, United States) (1977), 197(4310), 1289-90

CODEN: SCIEAS; ISSN: 0036-8075

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Polyribonucleosinic-polyribocytidylic acid stabilized with poly-L-lysine and carboxymethylcellulose [poly (ICLC)] enhances the antibody response in rhesus monkeys immunized with swine influenza virus subunit vaccine. Monkeys given the vaccine-adjuvant combination had earlier and higher titers by 14 days compared to those that received vaccine alone. The potentiation of the antibody response of young monkeys given a split-virus vaccine in combination with poly (ICLC) suggests that this vaccine-adjuvant combination may similarly provide a potentially useful alternative approach to the immunization of pediatric and young adult age groups against swine influenza.

L3 ANSWER 25 OF 25 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1977:69998 CAPLUS

DOCUMENT NUMBER: 86:69998

TITLE: Interferon induction in cynomolgus and rhesus monkeys after repeated doses of a modified polyribonucleosinic-polyribocytidylic acid complex

AUTHOR(S): Sammons, M. L.; Stephen, E. L.; Levy, H. B.; Baron, S.; Hilmas, D. E.

CORPORATE SOURCE: U. S. Army Med. Res. Inst. Infect. Dis., Fort Detrick, Frederick, MD, USA

SOURCE: Antimicrobial Agents and Chemotherapy (1977), 11(1), 80-3

CODEN: AMACCQ; ISSN: 0066-4804

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Serum interferon activity was determined in 12 cynomolgus and 12 rhesus monkeys injected i.v. once daily for 10 days with 0.1-6.0 mg/kg of a stabilized poly(ribonucleosinic acid)-poly(ribocytidylic acid) complex, composed of poly(ribonucleosinic acid)-poly(ribocytidylic acid), poly-L-lysine, and carboxymethylcellulose [poly (ICLC)]. Interferon activity was detected 2 h after the first

injection, with maximum activity occurring 8 h after the second injection. A period of hyporesponsiveness occurred after the third injection of poly(ICLC) in all monkeys and lasted until the sixth injection in the rhesus monkeys, when interferon activity again became more elevated. The delayed rebound was not as apparent in cynomolgus monkeys. Rhesus monkeys injected with 6 mg/kg did not exhibit serious side effects.

=> file stng

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

91.41

91.62

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-19.50

-19.50

FILE 'STNGUIDE' ENTERED AT 18:48:41 ON 11 JUL 2007

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jul 6, 2007 (20070706/UP).

=> d his

(FILE 'HOME' ENTERED AT 18:43:07 ON 11 JUL 2007)

FILE 'CAPLUS' ENTERED AT 18:43:17 ON 11 JUL 2007

L1 94 S (POLY (5A) ICLC)

L2 2 L1 AND PREP/RL

E LYSINE+ALL/CT

E CARBOXYMETHYLCELLULOSE+ALL/CT

L3 25 S L1 AND LYSINE AND (CARBOXYMETHYLCELLULOSE OR "CARBOXYMETHYL C

FILE 'STNGUIDE' ENTERED AT 18:48:41 ON 11 JUL 2007

=>